

**Ordering Physician:**  
Precision Analytical

**DOB:** 1967-08-09  
**Age:** 50  
**Gender:** Male



**Collection Times:**  
2017-08-09 06:01AM (S)  
2017-08-09 06:31AM (S)  
2017-08-09 07:01AM (S)  
2017-08-09 05:01PM (S)  
2017-08-09 10:01PM (S)  
2017-08-09 06:01AM (U)  
2017-08-09 08:01AM (U)  
2017-08-09 05:01PM (U)  
2017-08-09 10:01PM (U)

## Hormone Testing Summary

### Key (how to read the results):




### Sex Hormones



### Testosterone

Age	Range
18-25	50-115
26-40	40-95
41-60	30-80
>60	25-60




### Adrenal Hormones See pages 4 and 5 for a more complete breakdown of adrenal hormones



Free cortisol best reflects tissue levels. Metabolized cortisol best reflects total cortisol production.

### Total DHEA Production

Age	Range
20-39	3000-5500
40-60	2000-4000
>60	1000-2500



cortisol  
metabolism

The following videos (which can also be found on the website under the listed names along with others) may aid your understanding:  
[DUTCH Plus Overview](#) (quick overview) [Estrogen Tutorial](#) [Male Androgen Tutorial](#) [Cortisol/CAR Tutorial](#)

**PLEASE BE SURE TO READ BELOW FOR ANY SPECIFIC LAB COMMENTS. More detailed comments can be found on page 8.**

- The patient collected an "Insomnia" salivary sample in the middle of the night. The cortisol result for this sample was 2.10ng/mL (expected range 0-0.9). Please see page 4 for cortisol and cortisone results for this sample.  
The Cortisol Awakening Response (CAR) was 2.99ng/mL (expected range 1.5-4.0) or 69.2% (range 50-160%). See page 5 for more details.



**Accession # 00280397**  
 Male Sample Report  
 123 A Street  
 Sometown, CA 90266



**Sex Hormones and Metabolites**

**Ordering Physician:**  
 Precision Analytical

**DOB:** 1967-08-09  
**Age:** 50  
**Gender:** Male

**Collection Times:**  
 2017-08-09 06:01AM (S)  
 2017-08-09 06:31AM (S)  
 2017-08-09 07:01AM (S)  
 2017-08-09 05:01PM (S)  
 2017-08-09 10:01PM (S)  
 2017-08-09 06:01AM (U)  
 2017-08-09 08:01AM (U)  
 2017-08-09 05:01PM (U)  
 2017-08-09 10:01PM (U)

Category	Test	Result	Units	Normal Range
<b>Progesterone Metabolites (Urine)</b>				
	b-Pregnanediol	Low end of range	110.0	ng/mg 75 - 400
	a-Pregnanediol	Low end of range	40.0	ng/mg 20 - 130
<b>Estrogens and Metabolites (Urine)</b>				
	Estrone(E1)	High end of range	15.3	ng/mg 4 - 16
	Estradiol(E2)	Above range	2.4	ng/mg 0.5 - 2.2
	Estriol(E3)	Within range	4.2	ng/mg 2 - 8
	2-OH-E1	Within range	3.73	ng/mg 0 - 5.9
	4-OH-E1	Above range	1.2	ng/mg 0 - 0.8
	16-OH-E1	Within range	0.7	ng/mg 0 - 1.2
	2-Methoxy-E1	Within range	2.1	ng/mg 0 - 2.8
	2-OH-E2	Above range	0.61	ng/mg 0 - 0.6
	4-OH-E2	Within range	0.1	ng/mg 0 - 0.3
	2-Methoxy-E2	Within range	0.3	ng/mg 0 - 0.8
	Total Estrogen	High end of range	30.34	ng/mg 10 - 34
<b>Androgens and Metabolites (Urine)</b>				
	DHEA-S	Low end of range	95.0	ng/mg 30 - 1500
	Androsterone	Low end of range	835.0	ng/mg 500 - 3000
	Etiocholanolone	Below range	387.0	ng/mg 400 - 1500
	Testosterone	Below range	21.6	ng/mg 25 - 115
	5a-DHT	Low end of range	8.2	ng/mg 5 - 25
	5a-Androstanediol	Low end of range	52.0	ng/mg 30 - 250
	5b-Androstanediol	Low end of range	46.0	ng/mg 40 - 250
	Epi-Testosterone	Low end of range	38.1	ng/mg 25 - 115

Hormone metabolite results from the previous page are presented here as they are found in the steroid cascade. See the Provider Comments for more information on how to read the results.

**Androgens**


**Age-Dependent Ranges**

Age	DHEA-S
20-39	150-1500
40-60	60-800
>60	30-300



Etiocholanolone		Androsterone	
20-39	800-1500	20-39	1500-3000
40-60	600-1200	40-60	1000-2000
>60	400-1000	>60	500-1000

5β-androstanediol		5α-androstanediol	
20-39	70-250	20-39	60-250
40-60	55-210	40-60	50-180
>60	40-150	>60	30-130

Testosterone		5α-DHT	
18-25	50-115	20-39	9-25
26-40	40-95	40-60	7-20
41-60	30-80	>60	5-16
>60	25-60		



**Estrogens**





**Accession # 00280397**

Male Sample Report  
123 A Street  
Somertown, CA 90266






**Adrenal**

**Ordering Physician:**  
Precision Analytical

**DOB:** 1967-08-09  
**Age:** 50  
**Gender:** Male

**Collection Times:**  
2017-08-09 06:01AM (S)  
2017-08-09 06:31AM (S)  
2017-08-09 07:01AM (S)  
2017-08-09 05:01PM (S)  
2017-08-09 10:01PM (S)  
2017-08-09 01:31AM (S\*)  
2017-08-09 06:01AM (U)  
2017-08-09 08:01AM (U)  
2017-08-09 05:01PM (U)  
2017-08-09 10:01PM (U)

Category	Test	Result	Units	Normal Range
<b>Free Cortisol and Cortisone (Saliva)</b>				
	Saliva Cortisol - Waking (W)	High end of range	4.32	ng/mL 1.6 - 4.6
	Saliva Cortisol - W+30 min.	High end of range	7.31	ng/mL 3.7 - 8.2
	Saliva Cortisol - W+60 min.	Within range	4.33	ng/mL 2.3 - 5.3
	Saliva Cortisol - Afternoon	Within range	1.2	ng/mL 0.4 - 1.5
	Saliva Cortisol - Night	Within range	0.32	ng/mL 0 - 0.9
	Saliva Cortisone - Waking (W)	High end of range	13.16	ng/mL 6.8 - 14.5
	Saliva Cortisone - W+30 min.	Above range	25.57	ng/mL 12.4 - 19.4
	Saliva Cortisone - W+60 min.	Above range	16.9	ng/mL 9.4 - 15.3
	Saliva Cortisone - Afternoon	Above range	7.98	ng/mL 2 - 7.1
	Saliva Cortisone - Night	Within range	2.02	ng/mL 0 - 4.8
	Saliva Cortisol Total	High end of range	17.48	ng/mL 9.6 - 19.3
	Saliva Cortisone Total	Above range	65.63	ng/mL 36 - 55
<b>Creatinine (Urine)</b>				
	Creatinine A (Waking)	Within range	0.45	mg/ml 0.3 - 3
	Creatinine B (Morning)	Within range	0.41	mg/ml 0.3 - 3
	Creatinine C (Afternoon)	Within range	0.9	mg/ml 0.3 - 3
	Creatinine D (Night)	Within range	0.88	mg/ml 0.3 - 3
<b>Cortisol Metabolites and DHEA-S (Urine)</b>				
	a-Tetrahydrocortisol (a-THF)	Within range	450.0	ng/mg 175 - 700
	b-Tetrahydrocortisol (b-THF)	Within range	2800.0	ng/mg 1750 - 4000
	b-Tetrahydrocortisone (b-THE)	Above range	6000.0	ng/mg 2350 - 5800
	Metabolized Cortisol (THF+THE)	High end of range	9250.0	ng/mg 4550 - 10000
	DHEA-S	Low end of range	95.0	ng/mg 30 - 1500
<b>Additional Cortisol and Cortisone (Saliva)</b>				
	* Saliva Cortisol - Insomnia	Above range	2.1	ng/mL 0 - 0.9
	* Saliva Cortisone - Insomnia	Above range	10.4	ng/mL 0 - 4.8



- The patient submitted an Insomnia salivary sample. The cortisol result for this sample was 2.10ng/mL (expected range 0-0.9) The cortisone result for this sample was 10.4 ng/mL (expected range 0-4.8)

The Cortisol Awakening Response (CAR) is the rise in salivary cortisol between the waking sample and the sample collected 30 (as well as 60) minutes later. This "awakening response" is essentially a "mini stress test" and is a useful measurement in addition to the overall up-and-down (diurnal) pattern of free cortisol throughout the day. **This patient shows a waking cortisol of 4.32 and an increase to 7.3 after 30.0 minutes. This is an increase of 2.99ng/mL or 69.2%.** Expected increases differ depending on the methods used. Preliminary research shows that 50-160% or 1.5-4.0ng/mL increases are common with samples collected 30 minutes after waking. These guidelines are considered research only. **This patient shows a salivary cortisol of 4.33 measured 60 minutes after waking. This is an increase of 0.01ng/mL or 0.23% compared to the waking sample.** To date, data suggests that expected results may be 0-70%, and this guideline is considered for research only.



**Accession # 00280397**  
 Male Sample Report  
 123 A Street  
 Sometown, CA 90266



**Organic Acid Tests (OATs)**

**Ordering Physician:**  
 Precision Analytical

**DOB:** 1967-08-09  
**Age:** 50  
**Gender:** Male

**Collection Times:**  
 2017-08-09 06:01AM (S)  
 2017-08-09 06:31AM (S)  
 2017-08-09 07:01AM (S)  
 2017-08-09 05:01PM (S)  
 2017-08-09 10:01PM (S)  
 2017-08-09 06:01AM (U)  
 2017-08-09 08:01AM (U)  
 2017-08-09 05:01PM (U)  
 2017-08-09 10:01PM (U)

Category	Test	Result	Units	Normal Range
<b>Nutritional Organic Acids</b>				
Vitamin B12 Marker (may be deficient if high) - (Urine)				
	Methylmalonate (MMA)	Within range	1.2 ug/mg	0 - 3
Vitamin B6 Markers (may be deficient if high) - (Urine)				
	Xanthurenate	Above range	6.8 ug/mg	0 - 2.1
	Kynurenate	Above range	30.1 ug/mg	0 - 9.3
Glutathione Marker (may be deficient if low or high) - (Urine)				
	Pyroglutamate	Below range	23.2 ug/mg	43 - 85
<b>Neurotransmitter Metabolites</b>				
Dopamine Metabolite - (Urine)				
	Homovanillate (HVA)	Low end of range	5.6 ug/mg	4.8 - 19
Norepinephrine/Epinephrine Metabolite - (Urine)				
	Vanilmandelate (VMA)	Within range	4.8 ug/mg	2.8 - 8
Melatonin (*measured as 6-OH-Melatonin-Sulfate) - (Urine)				
	Melatonin* (Waking)	Within range	42.4 ng/mg	10 - 85
Oxidative Stress / DNA Damage, measured as 8-Hydroxy-2-deoxyguanosine (8-OHdG) - (Urine)				
	8-OHdG (Waking)	Within range	4.0 ng/mg	0 - 8.8





## References

- Handen, K., et al., Potential protective effect on key steroidogenesis and metabolic enzymes and sperm abnormalities by fenugreek seeds in tests on epididymis of surviving diabetic rats. *Arch Physiol Biochem*, 2010, **116**(3): p. 146-55.
- Simonian, M.H., ACTH and thyroid hormone regulation of 3 beta-hydroxysteroid dehydrogenase activity in human fetal adrenal cortex cells. *J Steroid Biochem*, 1986, **25**(6): p. 1001-6.
- Kaiki, E.M., et al., Distribution of steroidogenic enzymes involved in androgen synthesis in polycystic ovaries: an immunohistochemical study. *Mol Hum Reprod*, 2000, **6**(5): p. 443-7.
- Deluca, D., et al., Inhibition of 17beta-hydroxysteroid dehydrogenase by phytoestrogens: comparison with other steroid metabolizing enzymes. *J Steroid Biochem Mol Biol*, 2005, **93**(2-3): p. 285-92.
- Zhang, S., et al., Endocrine disruptors of inhibiting testicular 3beta-hydroxysteroid dehydrogenase. *Chem Biol Interact*, 2019, **303**: p. 50-57.
- Tomlinson, J.W., et al., Impaired glucose tolerance and insulin resistance are associated with increased adipose 17beta-hydroxysteroid dehydrogenase type 1 expression and elevated hepatic glyceraldehyde kinase activity. *Diabetes*, 2008, **57**(11): p. 2652-60.
- Stomati, M., et al., Six-month oral dehydroepiandrosterone supplementation in early and late postmenopausal. *Gynecol Endocrinol*, 2010, **14**(5): p. 342-43.
- Tsichorzidou, T., J.W. Honour, and G.S. Conway, Altered cortisol metabolism in polycystic ovary syndrome: insulin enhances 5alpha-reduction but not the elevated adrenal steroid production rates. *Clin Endocrinol (Oxf)*, 2003, **58**(2): p. 590-7.
- Prager, N., et al., A randomized, double-blind, placebo-controlled trial to determine the effectiveness of bioassayed derived inhibitors of 5-alpha-reductase in the treatment of androgenetic alopecia. *J Altern Complement Med*, 2002, **8**(2): p. 143-52.
- Fujita, R., et al., Anti-androgen activities of *Gonoloma luridum*. *J Ethnopharmacol*, 2005, **102**(1): p. 107-12.
- Moradi, H.R., et al., The histological and histochemical effects of Urtica dioica extract on rat's prostate hyperplasia. *Vet Res Forum*, 2015, **6**(1): p. 23-9.
- Will, T., et al., Pycnem africanum for benign prostatic hyperplasia. *Cochrane Database Syst Rev*, 2002(1): p. CD001044.
- Azzouzi, F., et al., The 5-alpha-reductase isozyme family: a review of basic biology and their role in human diseases. *Am J Urol*, 2012, **202**: p. 503-21.
- Westerkjøl, L., et al., Body fat distribution and cortisol metabolism in healthy men: enhanced 5beta-reductase and lower cortisol: cortone metabolite ratios in men with fatty liver. *J Clin Endocrinol Metab*, 2003, **88**(10): p. 4924-31.
- Gambineri, A., et al., Increased clearance of cortisol by 5beta-reductase in a subgroup of women with adrenal hyperandrogenism in polycystic ovary syndrome. *J Endocrinol Invest*, 2003, **2**(2): p. 210-8.
- Ojima, M., et al., The inhibitory effects of glycyrrhizin and glycyrrhetic acid on the metabolism of cortisol and prednisolone-in vivo and in vitro studies. *Nihon Nishyuppo Gakkaishi*, 1990, **66**(5): p. 584-96.
- Dupe, S., et al., 11beta-hydroxysteroid dehydrogenase types 1 and 2 activity in subcutaneous adipose tissue in humans: implications in obesity and diabetes. *Clin Endocrinol (Oxf)*, 2015, **100**(1): p. E70-6.
- Esteva, C.L., et al., Proinflammatory cytokine induction of 11beta-hydroxysteroid dehydrogenase type 1 (11beta-HSD1) in human adipocytes is mediated by MKK, CREBP, and NF-kappaB. *J Clin Endocrinol Metab*, 2014, **99**(1): p. E160-8.
- Chapman, K., W. Holmes, and A. Seckl, 11beta-hydroxysteroid dehydrogenase: intracellular gatekeepers of tissue glucocorticoid action. *Physiol Rev*, 2013, **93**(3): p. 1139-206.
- Hinzpeter, J., et al., Green tea and one of its constituents, Epigallocatechin-3-gallate, are potent inhibitors of human 11beta-hydroxysteroid dehydrogenase type 1. *PLoS One*, 2014, **9**(1): p. e84468.
- Hu, G.X., et al., Curcumin is a potent and selective inhibitor of 11beta-hydroxysteroid dehydrogenase: 1. Improving lipid profiles in high-fat-diet-treated rats. *PLoS One*, 2013, **8**(3): p. e69376.
- Amason, A.G., et al., Coffee inhibits the reactivation of glucocorticoids by 11beta-hydroxysteroid dehydrogenase type 1: a glucocorticoid competitor in the anti-diabetic action of coffee? *FEBS Lett*, 2006, **580**(17): p. 4081-5.
- Jothib Richard, E., et al., Anti-stress Activity of Oidrum sanctum: Possible Effects on Hypothalamic-Primary-Adrenal Axis. *Phytother Res*, 2016, **30**(5): p. 805-14.
- Blum, A., et al., Monomelic charantia extract, a herbal remedy for type 2 diabetes, contains a specific 11beta-hydroxysteroid dehydrogenase type 1 inhibitor. *J Steroid Biochem Mol Biol*, 2012, **128**(1-2): p. 51-5.
- Hoshino, M., et al., Comprehensive study of urinary cortisol metabolites in hypertrophic and hypothyroid patients. *Clin Endocrinol (Oxf)*, 2005, **64**(1): p. 37-45.
- Tanjanya, M., K. Horita, and Y. Iano, Urinary cortisol metabolites in the assessment of peripheral thyroid hormone action: application for diagnosis of resistance to thyroid hormone. *Thyroid*, 1993, **3**(3): p. 229-33.
- Uehama, H., et al., Decreased steroidogenic enzyme 17,20-lyase and increased 17alpha-hydroxysteroid activity in type 2 diabetes mellitus. *Eur J Endocrinol*, 2002, **146**(3): p. 375-80.
- Nestler, J.E. and D.J. Jakubowicz, Decreases in ovarian cytochrome P450c17 alpha activity and serum free testosterone after reduction of insulin secretion in polycystic ovary syndrome. *NIH J Med*, 1996, **335**(9): p. 617-23.
- Eggerhaas, D., et al., The influence of ketoconazole on human adrenal steroidogenesis: incubation studies with tissue slices. *Clin Endocrinol (Oxf)*, 1991, **35**(2): p. 143-8.
- Kossov, D.C. and H.D. Cobly, Dose-dependent actions of spironolactone on the inner and outer zones of the guinea pig adrenal cortex. *Pharmacology*, 1992, **45**(1): p. 27-33.
- Hasegawa, E., et al., Effect of polyphenols on production of steroid hormones from human adrenocortical NC-H295R cells. *Biol Pharm Bull*, 2013, **36**(2): p. 228-37.
- Mari, T., et al., Reversal of inhibits androgen production of human adrenocortical H295R cells by polyphenols. *PLoS One*, 2012, **7**(12): p. e4174224.
- Andric, S.A., et al., Acute effects of lipoic acid on testosterone production and free testosterone in rat testicular steroidogenesis. *Environ Health Perspect*, 2000, **108**(10): p. 955-9.
- Kim, S.H., et al., Body Fat Mass is Associated with Ratio of Steroid Metabolites Reflecting 17,20-OH Activity in Prepubertal Girls. *J Clin Endocrinol Metab*, 2016, **101**(12): p. 4653-4660.
- Armani, D., G. Borroni, and M. Palermo, Reduction of serum testosterone in men by fenugreek. *Eur J Med*, 1995, **34**(11): p. 1158.
- Armani, D., et al., Licorice reduces serum testosterone in healthy women. *Steroids*, 2004, **69**(1-12): p. 753-6.
- Seravali, P. and A. Lobo, The effects of spironolactone on adrenal steroidogenesis in hirsute women. *Fertil Steril*, 1985, **44**(5): p. 595-9.
- Ayub, M. and M.J. Level, Inhibition of human adrenal steroidogenic enzymes in vitro by imidazole drugs including ketoconazole. *J Steroid Biochem*, 1989, **32**(4): p. 515-24.
- Wang, X., et al., Suppression of rat and human androgen biosynthesis enzymes by adenosine: Possible use for the treatment of prostate cancer. *Fitoterapia*, 2016, **111**: p. 66-72.
- Hu, T., et al., Sporn acid tissue activation by uchi ameliorates polycystic ovary syndrome in rat. *Nat Biotechnol*, 2017, **47**: p. 21-28.
- Sarkola, T., et al., Acute effect of actoin on androgens in premenopausal women. *Alcohol Alcohol*, 2000, **35**(1): p. 84-90.
- Corbould, A.M., et al., The effect of obesity on the ratio of type 3 17beta-hydroxysteroid dehydrogenase mRNA to cytochrome P450 aromatase mRNA in subcutaneous abdominal and intra-abdominal adipose tissue of women. *Int J Obes Relat Metab Disord*, 2002, **26**(2): p. 165-75.
- Kazemian, A., et al., Human 17beta-hydroxysteroid dehydrogenase type 5 is inhibited by urinary flavonoids. *Adv Exp Med Biol*, 2002, **505**: p. 151-61.
- Le Bail, C., et al., Effects of phytoestrogens on aromatase, 5beta-reductase and 17beta-hydroxysteroid dehydrogenase activities and human breast cancer cells. *Life Sci*, 2000, **66**(4): p. 281-91.
- Adankou, S.O. and E.O. Feronibi, Quercetin ameliorates strazhen-induced changes in the testicular function of rats. *Toxicol Ind Health*, 2016, **32**(7): p. 1278-85.
- Gard, C. and K.A. Brown, Obesity and breast cancer - Role of estrogens and the molecular underpinnings of aromatase regulation in breast adipose tissue. *Mol Cell Endocrinol*, 2018, **466**: p. 15-30.
- Randolph, J.F., et al., The effect of insulin on aromatase activity in isolated human endometrial glands and stroma. *Am J Obstet Gynecol*, 1987, **157**(6): p. 1534-9.
- Watanabe, M. and S. Nakajin, Forskolin up-regulates aromatase (CYP19) activity and gene transcripts in the human adrenocortical carcinoma cell line H295R. *J Endocrinol*, 2004, **180**(1): p. 125-33.
- Sanderston, J., et al., Induction and inhibition of aromatase (CYP19) activity by natural and synthetic flavonoid compounds in H295R human adrenocortical carcinoma cells. *Toxicol Sci*, 2004, **82**(1): p. 70-9.
- Takeuchi, T., et al., Effect of apigenin, glycyrrhizin and glycyrrhetic acid on ovarian androgen production. *Am J Clin Med*, 1991, **1**(6): p. 32-8.
- Holloway, A.C., et al., Arazine-induced changes in aromatase activity in estrogen sensitive target tissues. *J Appl Toxicol*, 2008, **28**(3): p. 260-70.
- Leppart, E.D., Modulation of Aromatase by Phytoestrogens. *Enzyme*, 2015, **2015**: p. 594655.
- Novaes, M.R., et al., The effects of dietary supplementation with Agriolite mushrooms and other medicinal fungi on breast cancer: evidence based medicine. *Clinics (Sao Paulo)*, 2011, **66**(12): p. 2133-9.
- Saoh, K., et al., Inhibition of aromatase activity by green tea extract catechins and their endocrinological effects of oral administration in rats. *Food Chem Toxicol*, 2002, **40**(7): p. 925-33.
- Eng, E.T., et al., Suppression of estrogen biosynthesis by procyanidin dimer in red wine and grape seeds. *Cancer Res*, 2003, **63**(23): p. 8516-22.
- Chen, J., et al., The correlation of aromatase activity and obesity in women with or without polycystic ovary syndrome. *J Ovarian Res*, 2015, **8**: p. 11.
- Ayub, M. and M.J. Level, The inhibition of human prostatic aromatase activity by imidazole drugs including ketoconazole and 4-hydroxyandrostenedione. *Biochem Pharmacol*, 1990, **40**(7): p. 1569-75.
- Ren, S., et al., Dual effect of melatonin on growth inhibition and steroid production in breast cancer cells. *Int J Med Biol*, 2015, **35**(4): p. 1088-94.
- Richard, S., et al., Differential effects of glyphosate and roundup on human placental cells and aromatase. *Environ Health Perspect*, 2005, **113**(6): p. 716-20.
- Hogdes, M.E. and D.M. Wilchik, Modulation of Metabolic Deoxycation Pathways Using Foods and Food-Derived Compounds: A Scientific Review with Clinical Application. *Nutr Metab*, 2015, **2015**: p. 676889.
- Michonvici, J.J., H. Adlercreutz, and H.L. Bradlow, Changes in levels of urinary estrogen metabolites after oral intake 3-cachino treatment in humans. *J Natl Cancer Inst*, 1997, **89**(10): p. 718-23.
- Sowers, M.R., et al., Selected diet and lifestyle factors are associated with estrogen metabolites in a multicenteral population of women. *J Nutr*, 2006, **136**(6): p. 1588-95.
- Lu, J.J., et al., Increased urinary excretion of 2-hydroxyestrogen but not 16alpha-hydroxyestrogen in premenopausal women during a soy diet containing isoflavones. *Cancer Res*, 2000, **60**(5): p. 1299-305.
- Chen, H.Y., et al., The combined effects of garlic oil and fish oil on the hepatic antioxidant and drug-metabolizing enzymes of rats. *Br J Nutr*, 2003, **98**(2): p. 188-200.
- DeVease, P., et al., Induction of cytochrome P450 and/or detoxication enzymes by various extracts of rosemary: description of specific patterns. *Food Chem Toxicol*, 2001, **39**(9): p. 907-18.
- Michonvici, J.J. and K.A. Galbraith, Effects of exogenous thyroxine on C-6 and C-16 alpha hydroxylations of estradiol in humans. *Steroids*, 1990, **55**(1): p. 22-6.
- Peters, L.P. and R.W. Teal, Effect of high sucrose diet on cytochrome P450 1A and heterocyclic amine mutagenesis. *Anticancer Res*, 2003, **23**(1A): p. 399-403.
- Mahabir, S., et al., Effects of low-to-moderate alcohol supplementation on urinary estrogen metabolites in postmenopausal women in a controlled feeding study. *Cancer Med*, 2017, **6**(10): p. 2419-2423.
- Luzarasa, B., et al., Resveratrol and its methoxy derivatives modulate the expression of estrogen metabolism enzymes in breast epithelial cells by Akt down-regulation. *Mol Cell Biochem*, 2017, **425**(1-2): p. 169-179.
- Smerdova, L., et al., Up-regulation of CYP1B1 expression by inflammatory cytokines is mediated by the p38 MAP kinase signal transduction pathway. *Carcinogenesis*, 2014, **35**(11): p. 2534-43.
- Li, M.X., et al., Estrogen receptor alpha promotes smoking carcinogen-induced lung carcinogenesis via cytochrome P450 1B1. *Mol Med (Biol)*, 2015, **93**(11): p. 1221-33.
- Jaramillo, I.C., et al., Effects of (1a) components and combustion particle physicochemical properties on toxicological responses of lung cells. *Environ Sci Health A: Toxic Hazard Subst Environ Res*, 2018, **53**(4): p. 295-309.
- Doodnar, H., M.D. Burke, and R.T. Mayer, Bioflavonoids: selective substrates and inhibitors for cytochrome P450 CYP1A and CYP1B1. *Toxicology*, 2000, **144**(1-3): p. 31-8.
- Whitten, D.L., et al., The effect of St John's wort extracts on CYP3A: a systematic review of prospective clinical trials. *Br J Clin Pharmacol*, 2006, **62**(5): p. 512-26.
- Bhargava, H.L., et al., Effects of pesticides on the ratio of 16 alpha beta2-hydroxysterone: a biological marker of breast cancer risk. *Environ Health Perspect*, 1995, **103**(Suppl 7): p. 147-50.
- Luckert, C., et al., Polycyclic aromatic hydrocarbons stimulate human CYP3A4 promoter activity via PKR. *Toxicol Lett*, 2013, **222**(2): p. 180-8.
- Wu, W.H., et al., Estrogenic effect of yam ingestion in healthy postmenopausal women. *Am J Clin Nutr*, 2005, **24**(4): p. 235-43.
- Dresser, G.K., et al., Evaluation of peppermint oil and acetyl salicylate as inhibitors of cytochrome P4503A4 activity in vitro and in vivo. *Clin Pharmacol Ther*, 2002, **72**(3): p. 247-55.
- Niwa, T., Y. Imagawa, and H. Yamazaki, Drug interactions between nine antiulcer agents and drugs metabolized by human cytochromes P450. *Curr Drug Metab*, 2014, **15**(7): p. 651-79.
- Jiang, H., et al., Human catechol-O-methyltransferase down-regulation by estradiol. *Neuropharmacology*, 2003, **45**(7): p. 1011-8.
- Ho, P.W., et al., Effects of plasticizers and related compounds on the expression of the soluble form of catechol-O-methyltransferase in MCF-7 cells. *Curr Drug Metab*, 2008, **9**(4): p. 276-9.
- Blum, K., et al., Manipulation of catechol-O-methyltransferase (COMT) activity to influence the attenuation of substance seeking behavior: a subtype of Reward Deficiency Syndrome (RDS) is dependent upon gene polymorphisms: a hypothesis. *Mol Hypothesis*, 2007, **6**(9): p. 1054-60.
- van Duursen, M.B., et al., Phytochemicals inhibit catechol-O-methyltransferase activity in systemic fractions from healthy human mammary tissues: implications for catechol estrogen-induced DNA damage. *Toxicol Sci*, 2004, **81**(2): p. 316-24.
- Sehmi, A.O., et al., St. John's wort ameliorate 2,4,6-trinitrobenzenesulfonic acid colitis of rats through the induction of pregnane X receptor and P-glycoproteins. *Physiol Pharmacol*, 2015, **66**(2): p. 203-14.
- Pastuss, J.M., et al., Deventanethasone induces pregnane X receptor and rethoid X receptor-alpha expression in human hepatocytes: synergistic increase of CYP3A4 induction by pregnane X receptor activators. *Mol Pharmacol*, 2000, **58**(2): p. 361-72.
- Zhou, H. and P.B. Hylonen, Bile acids are nutrient signaling hormones. *Steroids*, 2014, **86**: p. 62-8.
- Ding, X. and L.L. Staudinger, Induction of drug metabolism by farnesol: the role of the pregnane X receptor and the protein kinase A signal transduction pathway. *J Pharmacol Exp Ther*, 2005, **312**(2): p. 889-95.
- Muelier, J.W., et al., The Regulation of Steroid Action by Sulfation and Deacetylation. *Endocr Rev*, 2015, **36**(5): p. 526-63.
- Kim, M.S., et al., Suppression of Dihydro sulfotransferase (SULT2A1) during the acute phase response. *Am J Physiol Endocrinol Metab*, 2004, **287**(4): p. E731-8.
- Al-Dajali, E.A., et al., Lipoic acid and glycyrrhetic acid increase DHEA and deoxyandrosterone levels in vivo and in vitro by inhibiting adrenal SULT2A1 activity. *Mol Cell Endocrinol*, 2011, **336**(1-2): p. 102-9.


ref 021220




# Provider Notes

## How to read the DUTCH report

This report is not intended to treat, cure or diagnose any specific diseases. The graphic dutch dials in this report are intended for quick and easy evaluation of which hormones are out of range. Results below the left star are shaded yellow and are below range (left). Results between the stars and shaded green are within the reference range (middle). Results beyond the second star and shaded red are above the reference range (right). Some of these hormones also change with age, and the age-dependent ranges provided should also be considered.



In a few places on the graphical pages, you will see fan-style gauges. For sex hormones, you will see one for the balance between 5a/5b metabolism as well as methylation. For adrenal hormones, you will see one to represent the balance between cortisol and cortisone metabolites. These indexes simply look at the ratio of hormones for a preference. An average or "normal" ratio between the two metabolites (or groups of metabolites) will give a result in the middle (as shown here). If the ratio between the metabolites measured is "low" the gauge will lean to the left and similarly to the right if the ratio is higher than normal.



## Patient or Sample Comments

Throughout the provider comments you may find some comments specific to your situation or results. These comments will be found in this section or within another section as appropriate. Comments in other sections that are specific to your case will be in **bold**.

Note: The dates listed on the samples imply that they were older than our allowed 3 weeks when they were received. The instructions ask that patients freeze or refrigerate samples if they are to be held. If that is not the case, the free cortisol and cortisone levels may drop somewhat over time if the samples are too old. Other hormones tested are stable for more than 12 weeks at room temperature. Samples that are refrigerated or frozen are stable for months.

## Androgen Metabolism

When evaluating androgen levels, it is important to assess the following:

### • The status (low, normal or high?) of DHEA:

DHEA and androstenedione are made almost exclusively by the adrenal gland (although a smaller amount is made in the ovaries). These hormones appear in urine as DHEA-S (DHEA-Sulfate), androsterone and etiocholanolone. The best way to assess the total production of DHEA is to add up these three metabolites. This total can be seen on the first page of the DUTCH Complete (and DUTCH Plus). DHEA production decreases quite significantly with age. Age-dependent ranges can be seen on the graphical page of results.

**The Total DHEA Production (page 1) was about 1,317ng/mg which is within the overall range but is below the range for the patient's age-dependent range. This implies that the adrenal glands are not producing appropriate DHEA levels for the patient's age. Low DHEA is associated with depression, diabetes, heart disease, inflammation and immune disorders. It can be caused by hypothyroidism. It can cause fatigue, low mood and low libido. Supplementing DHEA in women often raises both testosterone and estrogen, which may or may not be desirable here. DHEA may increase with adaptogens such as maca and rhodiola, which improve overall adrenal output.**


### • The status (low, normal or high?) of testosterone:

The testes make most of the male's testosterone. Levels tend to be their highest at around 20 years of age and start to decline when men get into their 30's. Levels continue to drop as men age. Consider the appropriate age-dependent range for your patient. In older men, you can also consider the 18-25 year-old group to approximate what levels may have been when the patient was young and relatively healthy.

### • The metabolic preference for the 5a (5-alpha) or 5b (5-beta) pathway:

5a-reductase converts testosterone into 5a-DHT (DHT), which is even more potent (~3x) than testosterone. High levels of DHT can lead to symptoms associated with too much testosterone (thinning scalp hair, acne, etc.) and may also be associated with prostate issues in older men. Metabolites created down the 5b-pathway are significantly less androgenic than their 5a counterparts. In the examples below, the example on the left shows a patient with 5b-metabolism preference. A patient with a pattern like the example on the right may have high androgen symptoms even though testosterone is in the normal range because of the likely preference for turning a lot of his testosterone into DHT. The fan-style gauge below the hormones shows the 5a or 5b preference based on the balance between etiocholanolone (5b) and androsterone (5a) as well as 5a-androstanediol and 5b-androstanediol.

Example of how to read fan-style gauge for 5a-reductase activity:



You will also see levels of epi-testosterone, which is not androgenic like testosterone. It happens to be produced in about the same concentrations as testosterone (this is an approximate relationship). This can be helpful to assess testosterone therapy and rare cases where testosterone may have other complexities.

### Estrogen Metabolism

When evaluating estrogen levels, it is important to assess the following:

- **The status (low, normal or high?) of estrogen production:**

Levels of the primary estrogen, estradiol (the strongest estrogen), as well as "total estrogens" may be considered.

- **Phase I Metabolism:**

Estrogen is metabolized (primarily by the liver) down three phase I pathways. The 2-OH pathway is considered the safest because of the anti-cancer properties of 2-OH metabolites. Conversely, the 4-OH pathway is considered the most genotoxic as its metabolites can create reactive products that damage DNA. The third pathway, 16-OH creates the most estrogenic of the metabolites (although still considerably less estrogenic than estradiol) - 16-OH-E1.

When evaluating phase I metabolism, it may be important to look at the ratios of the three metabolites to see which pathways are preferred relative to one another. It may also be important to compare these metabolites to the levels of the parent hormones (E1, E2). If the ratios of the three metabolites are favorable but overall levels of metabolites are much lower than E1 and E2, this may imply sluggish phase I clearance of estrogens, which can contribute to high levels of E1 and E2.

The pie chart will assist you in comparing the three pathway options of phase I metabolism compared to what is "normal." 2-OH metabolism can be increased by using products containing D.I.M. or I-3-C. These compounds are found (or created from) in cruciferous vegetables and are known for promoting this pathway.

- **Methylation (part of Phase II Metabolism) of estrogens:**

After phase I metabolism, both 4-OH and 2-OH (not 16-OH) estrogens can be deactivated and eliminated by methylation. The methylation-activity index shows the patient's ratio of 2-Methoxy-E1 / 2-OH-E1 compared to what is expected. Low methylation can be caused by low levels of nutrients needed for methylation and/or genetic abnormalities (COMT, MTHFR). The COMT enzyme responsible for methylation requires magnesium and methyl donors. Deficiencies in folate or vitamin B6 or B12 can cause low levels of methyl donors. MTHFR genetic defects can make it more difficult for patients to make sufficient methyl donors. Genetic defects in COMT can make methylation poor even in the presence of adequate methyl donors.

Progesterone levels are of marginal value in men, although deficiency can be associated with some clinical conditions such as depression, fatigue, and low libido.

**Progesterone metabolites have limited relevance in male patients, but may be worth considering in some patients with abnormal results. In this case, both progesterone metabolites are within the normal range.**

### DUTCH Adrenal

The HPA-Axis refers to the communication and interaction between the hypothalamus (H) and pituitary (P) in the brain down to the adrenal glands (A) that sit on top of your kidneys. When a physical or psychological stressor occurs, the hypothalamus tells the pituitary to make ACTH, a hormone. ACTH stimulates the adrenal glands to make the stress hormone, cortisol and to a lesser extent DHEA and DHEA-S. Normally, the HPA-axis production follows a daily pattern in which cortisol rises rather rapidly in the first 10-30 minutes after waking (this is the C.A.R.) in order to help with energy, then gradually decreases throughout the day so that it is low at night for sleep. The cycle starts over the next morning. Abnormally high activity occurs in Cushing's Disease where the HPA-axis is hyper-stimulated causing cortisol to be elevated all day. The opposite is known as Addison's Disease, where cortisol is abnormally low because it is not made appropriately in response to ACTH's stimulation. These two conditions are somewhat rare. Examples of more common conditions related to less severely abnormal cortisol levels include fatigue, depression, insomnia, fibromyalgia, anxiety, inflammation and more.

Only a fraction of cortisol is "free" and bioactive. This fraction of cortisol is very important, but levels of metabolized cortisol

best represent overall production of cortisol therefore both should be taken into account to correctly assess adrenal function.

When evaluating cortisol levels, it is important to assess the following:

- **The overall up-and-down pattern of free cortisol throughout the day, looking for low and high levels:**

Abnormal results should be considered along with related symptoms.

- **The sum of the free cortisol as an expression of the overall tissue cortisol exposure:**

This total of five free cortisol measurements is the best way to assess the total of free cortisol throughout the day, but do be aware that it is heavily weighted towards the morning production since three of five measurements are made within the first hour of the day.

- **The total level of cortisol metabolites:**

We call this calculation "Metabolized Cortisol" which is the sum of a-THF, b-THF and b-THE. While free cortisol is the best assessment for tissue levels of cortisol, it only represents 1-3% of the total produced. The majority of cortisol results in a urine metabolite and the total of these metabolites best represents the total glandular output of cortisol for the day. When overall production is much higher than free cortisol levels, cortisol clearance may be increased (as seen in hyperthyroidism, obesity, etc.) The most common reason for sluggish cortisol clearance (assumed when free cortisol levels are much higher than metabolized cortisol) is low thyroid.

- **A potential preference for cortisol or cortisone (the inactive form):**

Looking at the comparison between the total for free cortisol and free cortisone is NOT the best indication of a person's preference for cortisol or cortisone. The saliva gland converts cortisol to cortisone in the local tissue. This localized conversion can be seen by comparing cortisol and cortisone levels. To see the patient's preference systemically, it is best to look at which *metabolite* predominates (THF or THE). This preference can be seen in the gauge below metabolized cortisol. This is known as the 11b-HSD index. The enzyme 11b-HSD II converts cortisol to cortisone in the kidneys, saliva gland and colon. 11b-HSD I is more active in the liver, fat cells and the periphery and is responsible for reactivating cortisone to cortisol. Both are then metabolized by 5a-reductase to become tetrahydrocortisol (THF) and tetrahydrocortisone (THE) respectively.

- **The Cortisol Awakening Response (CAR):**

The unique feature of the DUTCH Plus is the inclusion of the CAR assessment. The response to waking adds one more piece to HPA-axis function. In some cases overall levels of free cortisol may be normal, but the response to stress may be under or overactive. Reasons for a lower CAR might include: an underactive HPA Axis, excessive psychological burnout, seasonal affective disorder (SAD), sleep apnea or poor sleep in general, PTSD, and "chronic fatigue" patients. An elevated CAR can be a result of an over-reactive HPA axis, ongoing job-related stress (anticipatory stress for the day), glycemic dysregulation, pain (ie. waking with painful joints or a migraine), and general depression (not SAD). Scientific literature points to the magnitude of the morning cortisol increase as being connected to HPA-axis health whether the overall production of cortisol is low, normal or high.

**- The patient submitted an Insomnia salivary sample. The cortisol result for this sample was 2.10ng/mL. The cortisone result was 10.4 ng/mL. Ranges can be found in the table on the last page.**

### Nutritional Organic Acids

The following three organic acids are functional markers for vitamin deficiency. These compounds essentially back up in human biochemistry when a key nutrient is missing. These three metabolites have fairly straightforward interpretations. When the markers are elevated, it is likely that the patient's cellular levels of the related nutrient may be insufficient.

#### Methylmalonate (MMA)

Methylmalonate (also known as methylmalonic acid or MMA) is a functional marker of vitamin B12 (also known as cobalamin) deficiency. When cellular levels of B12 are low either from deficiency or due to a B12 transporter gene mutation, levels of MMA increase. This marker is considered superior to measuring serum B12 levels directly. A 2012 publication by Miller showed that 20% of those tested had a genetic defect in the protein that transports B12 to cells. These patients may have a functional B12 deficiency even if serum levels of B12 are normal.

If levels of MMA are elevated, it may be advisable to increase B12 consumption. Common foods high in B12 include beef liver, sardines, lamb, wild caught salmon, grass-fed beef, nutritional yeast and eggs. Vitamin B12 levels can also be increased through supplementation of B12 (taken as cobalamin, methylcobalamin, hydroxycobalamin, or adenosylcobalamin). Symptoms of a vitamin B12 deficiency include: fatigue, brain fog, memory problems, muscle weakness, unsteady gait, numbness, tingling, depression, migraines/headaches and low blood pressure.

#### Xanthurenate

Xanthurenate (also known as xanthurenic acid) and Kynurenate (kynurenic acid) are functional markers of vitamin B6 (also known as pyridoxine) deficiency. Vitamin B6 is a critical co-factor to over 100 important reactions that occur in the human body and is stored in the highest concentrations in muscle tissue. Tryptophan is readily converted to NAD by the liver. One of the steps in this pathway requires B6. When there is insufficient B6, xanthurenate is made instead. Kynurenate may also become elevated when patients are B6 deficient because of a different, possibly less B6 dependent pathway. The pathways leading to these biomarkers have other influences, so they will not always agree. When Xanthurenate is elevated, Kynurenate is also elevated about 1/3 of the time. When both are elevated, a B6 deficiency is likely more certain and more severe.

Not only is xanthurenate an indicator of a lack of B6, it is also harmful to the human body. It complexes with insulin and decreases insulin sensitivity. In fact, rats fed xanthurenate will actually develop diabetes because of the effects on insulin. If xanthurenate levels are elevated, B6 supplementation may be considered. Food high in B6 include turkey breast, grass-fed beef, pinto beans, avocado, pistachios, chicken, sesame and sunflower seeds.

While there is always some tryptophan going down the kynurenine pathway towards NAD (and possibly xanthurenate), this

process is up-regulated by inflammation, estrogen and cortisol. If levels of estrogen or cortisol are high, it may exacerbate xanthurenate elevations and increase the need for B6.

Xanthurenate can also bind to iron and create a complex that increases DNA oxidative damage resulting in higher 8-OHdG levels. If both markers are elevated, there is likely an antioxidant insufficiency.

**Xanthurenate and Kynurenate are both elevated in this case, so a vitamin B6 deficiency is likely and may be somewhat significant (since both markers are elevated). It is advisable to consider increasing vitamin B6 intake and to be aware of those things listed above that may induce a vitamin B6 deficiency.**

### **Pyroglutamate**

Pyroglutamate (also known as pyroglutamic acid) is a functional marker of glutathione deficiency. Pyroglutamate is a step in the production/recycling of glutathione. If the body cannot convert pyroglutamate forward, it will show up elevated in the urine. High pyroglutamate is an established marker for glutathione deficiency. Pyroglutamate in the urine can also be elevated with cheese consumption.

Glutathione is one of the most potent anti-oxidants in the human body. It is especially important in getting rid of toxins, including the reactive quinone species formed by 4-OH-E1 and 4-OH-E2. This reactive species can damage DNA if not detoxified by either methylation or glutathione.

Some have reported that low pyroglutamate may also be indicative of a need for glutathione; however, this is not established in the scientific literature.

### **Neurotransmitter Metabolites**

The neurotransmitters dopamine, norepinephrine and serotonin are important for human health. Measuring neurotransmitters directly (direct testing of serotonin, for example) is difficult because of their instability and their urinary measurements are controversial with respect to how well they reflect the body's levels of these neuro-hormones. Each of these three neurotransmitters can be assessed indirectly by measuring their urine metabolites. While these metabolites are not a perfect reflection of what's going on in the brain, the scientific literature does affirm their use for a good representation of overall levels of these neurotransmitters.

### **Homovanillate (HVA)**

Homovanillate (also known as HVA) is the primary metabolite of dopamine, a brain and adrenal neurotransmitter that comes from tyrosine (with BH4 and iron as co-factors) and goes on to create norepinephrine (noradrenaline) and epinephrine (adrenaline).

Low levels of HVA can be due to low levels of dopamine or poor conversion of dopamine to HVA. The latter may be due to insufficient levels of SAM, Magnesium, FAD and NAD which are needed to metabolize dopamine. Low circulating dopamine may be due to insufficient BH4, iron or tyrosine. It may also be seen when adrenal function is generally low. Low dopamine levels may be associated with addictions, cravings and pleasure seeking (to boost levels) in addition to sleepiness, impulsivity, tremors, less motivation, fatigue and low mood.

Elevated HVA may be caused by generally increased adrenal hormone output or because of a copper or vitamin C deficiency (which are needed for dopamine conversion to norepinephrine). Elevations may also be caused by a number of medications or supplements including: MAO inhibitors, quercetin, tyrosine, DL-phenylalanine (DLPA), L-dopa, macuna, dopamine medication (Levodopa, Sinemet, Methyl dopa), SNRI medication (Wellbutrin), tricyclic antidepressants, amphetamines, appetite suppressants, and caffeine. Bananas also contain dopamine. Elevated dopamine may be associated with loss of memory, insomnia, agitation, hyperactivity, mania, hyper-focus, high stress and anxiety as well as addictions, cravings and pleasure seeking (to maintain high levels).

When HVA is very high, consider if the previously discussed foods, supplements or medications may be the cause. Rarely, tumors associated with increased HVA may be present. In these cases, further testing is necessary for diagnosis. High HVA alone is not diagnostic of a tumor.

### **Vanilmandelate (VMA)**

Vanilmandelate (also known as VMA) is the primary metabolite of norepinephrine and epinephrine (adrenaline). The adrenal gland makes cortisol and DHEA as well as norepinephrine and epinephrine. When adrenal hormone output is generally low, VMA levels may be low. If HVA levels are significantly higher than VMA, there may be a conversion problem from dopamine to norepinephrine. This case can be caused by a copper or vitamin C deficiency. The enzymes COMT (methylation) and MAO are needed to make VMA from norepinephrine. If these enzymes are not working properly, VMA may be low when circulating norepinephrine and/or epinephrine are not low. Low levels of norepinephrine and epinephrine may be associated with addictions, cravings, fatigue, low blood pressure, low muscle tone, intolerance to exercise, depression, loss of alertness.

When the body is under physical or psychological stress, VMA levels may increase. Because dopamine gets converted to norepinephrine and ultimately to VMA, the list of medications and supplements that increase HVA may also increase VMA. Elevated levels may be associated with feeling stressed, aggression, violence, impatience, anxiety, panic, worry, insomnia, paranoia, increased tingling/burning, loss of memory, pain sensitivity, high blood pressure and heart palpitations.

If VMA and HVA are both extremely high, it may be necessary to rule out a neuroblastic tumor.

### **Melatonin (measured as 6-OHMS)**

Melatonin is not technically an adrenal or sex hormone however it is highly involved in the entire endocrine system. It is made in small amounts in the pineal gland in response to darkness and stimulated by Melanocyte Stimulating Hormone (MSH). A low MSH is associated with insomnia, an increased perception of pain, and mold exposure. Pineal melatonin (melatonin is also made in significant quantities in the gut) is associated with the circadian rhythm of all hormones (including female hormone release). It is also made in small amounts in the bone marrow, lymphocytes, epithelial cells and mast cells. Studies have shown that a urine sample collected upon waking has levels of 6-Hydroxymelatonin-sulfate (6-OHMS) that correlate well to the total levels of melatonin in blood samples taken continuously throughout the night. The DUTCH test uses

the waking sample only to test levels of melatonin production.

Low melatonin levels may be associated with insomnia, poor immune response, constipation, weight gain or increased appetite. Elevated melatonin is usually caused by ingestion of melatonin through melatonin supplementation or eating melatonin-containing foods. Elevated melatonin production that is problematic is rare, but levels can be higher in patients with Chronic Fatigue Syndrome and may be phase shifted (peaking later) in some forms of depression.

### **8-OHdG (8-Hydroxy-2-deoxyguanosine)**

8-OHdG (8-hydroxy-2-deoxyguanosine) results can be seen on page 6 of the DUTCH Complete (or DUTCH Plus) report. It is a marker for estimating DNA damage due to oxidative stress (ROS creation). 8-OHdG is considered pro-mutagenic as it is a biomarker for various cancer and degenerative disease initiation and promotion. It can be increased by chronic inflammation, increased cell turnover, chronic stress, hypertension, hyperglycemia/pre-diabetes/diabetes, kidney disease, IBD, chronic skin conditions (psoriasis/eczema), depression, atherosclerosis, chronic liver disease, Parkinson's (increasing levels with worsening stages), Diabetic neuropathy, COPD, bladder cancer, or insomnia. Studies have shown higher levels in patients with breast and prostate cancers. When levels are elevated it may be prudent to eliminate or reduce any causes and increase the consumption of antioxidant containing foods and/or supplements.

The reference range for 8-OHdG is a more aggressive range for Functional Medicine that puts the range limit at the 80th percentile for each gender. A classic range (average plus two standard deviations) would result in a range of 0-6ng/mg for women and 0-10ng/mg for men. Seeking out the cause of oxidative stress may be more crucial if results exceed these limits.



